

## **Virus self-assembly modeling**

***Per Linse and Ran Zang***

Physical Chemistry

Department of Chemistry

Lund University, P.O. Box 124

S-221 00 Lund, SWEDEN

per.linse@fkem1.lu.se

The structural organization of viruses is characterized by simplicity and economy. At a minimal level, the viral genome is composed of either single or double stranded RNA or DNA, surrounded by a spherical-like or cylindrical protein shell composed of many copies of almost identical proteins. The linear size of the virus genome can be tens of microns and is large compared to the diameter of capsid, usually in the range of tens of nanometers, so the genome must undergo a high degree of compaction during viral assembly. *In vivo*, (1) spherical-like viruses with double-stranded genome are made in two steps; first an empty capsid is formed and thereafter the genome is actively transferred into the capsid, whereas (2) spherical-like viruses with single-stranded genome are formed by a co-assembly of capsid subunits and the genome.

During the last decade, spherical-like viruses have been subjected to theoretical modeling using descriptions of different degree of details. After an introduction, results from molecular dynamic studies of a coarse-grain model describing the process of forming single-stranded viruses possessing icosahedral symmetry will be discussed. In particular, the influence of (i) net charge and charge distribution of capsid subunits and (ii) the topology of the genome on the virus assembly process will be highlighted. The fundamental understanding of the nature of viruses is essential for treating virus diseases as well as biomimetic inspired constructions of nanocarrier for medical and technical use.